



The status and future of the Lake Naivasha fishery, Kenya

Phil Hickley¹, Roland Bailey², David M. Harper, Rodrick Kundu³, Mucai Muchiri⁴,
Rick North¹ & Andy Taylor¹

University of Leicester, Leicester LE1 7RH, U.K.

¹Present address: National Coarse Fisheries Centre, Environment Agency, Arthur Drive,
Hoo Farm Industrial Estate, Worcester Road, Kidderminster, DY11 7RA, U.K.

²Life Sciences Department, King's College London, 150 Stamford Street, London SE1 9NN, U.K.

³Present address: Fisheries Department, P.O. Box 47066, Nairobi, Kenya

⁴Present address: Department of Fisheries, Moi University, P.O. Box 3900, Eldoret, Kenya

Key words: fishery yield, introduced species, *Oreochromis leucostictus*, *Tilapia zillii*, *Micropterus salmoides*

Abstract

Lake Naivasha is a freshwater lake situated in the eastern rift valley of Kenya. Only five species of fish are present, all of which have been introduced. They are *Oreochromis leucostictus*, *Tilapia zillii*, *Micropterus salmoides* (largemouth bass), *Barbus amphigramma* and *Poecilia reticulata* (guppy). The first three of these form the basis of an important gill net fishery and bass are also taken by rod and line for sport. *Barbus* are occasionally caught by dip net. Actual and potential yields for the Lake Naivasha fishery are discussed and the fishery is shown to be under-performing. The feeding regimes of the commercially exploited fish were examined in the context of available food supply, in particular the benthic fauna. Small bass depend heavily on *Micronecta* and large bass mostly take crayfish. Detritus predominates in the diet of *Oreochromis leucostictus* and *Tilapia zillii* but the former also eats algae and the latter, *Micronecta* and macrophyte. Various food resources, especially the benthos, appear to be under-utilized and so it is possible that further species could be introduced to enhance the commercial fishery.

Introduction

Lake Naivasha is a shallow (3–6 m) freshwater lake, approximately 160 km² in area, situated in the eastern rift valley of Kenya about 100 km north of Nairobi. It lies in a closed basin at an altitude of 1890 m above sea level, receives 90% of its water from the perennial River Malewa, and is subject to considerable fluctuations in water level. Dominant vegetation types are belts of papyrus (*Cyperus papyrus* L.) around the margins, stands of submerged macrophytes, of which the principal species is *Najas pectinata* (Parl.), and mats of floating plants comprising *Salvinia molesta* Mitch. and *Eichhornia crassipes* (Mart.). An overview of the lake and its changing ecology can be found in Harper et al. (1990).

Resulting from a probable history of occasional drying out, Lake Naivasha when first studied (c.1900) had only one species of fish present, the endemic

Aplocheilichthys antinorii (Vinc.) which was last recorded in 1962 (Elder et al., 1971). Since 1925, various fish introductions have been made, some successful and some not (Litterick et al., 1979; Muchiri & Hickley, 1991). Present today are *Oreochromis leucostictus* (Trewewas), *Tilapia zillii* (Gervais), *Micropterus salmoides* Lacépède (largemouth bass), *Barbus amphigramma* Blgr. and *Poecilia reticulata* Peters (guppy). Since 1959, the two tilapias and the bass have formed an important fishery (Muchiri & Hickley, 1991) with all three species being commercially exploited using gill nets and the bass also being taken by rod and line for sport.

The purpose of this paper is to relate the current status of the commercial fishery to information on fish feeding regimes and potential yields so as to identify whether or not there is scope for better management and utilisation of the resource.

Methods

Commercial catch data were obtained from the Fisheries Department of the Kenya Government. Fishermen use multifilament gill nets with a minimum stretched mesh size of 100 mm and are limited to a maximum of 10 nets of not more than 100 m in length. All catches must be landed at a single station near Naivasha town where fisheries personnel record the weight of the catch from each canoe. Data collection commenced in 1963. In addition, some *B. amphigramma* were taken downstream of a weir on the River Malewa.

Gill net catch statistics were fitted to a version of the Schaefer model (Ricker, 1975; Pauly, 1983): $C = a + bE$ where C = catch per unit effort, E = effort (with effort being the number of canoes licensed to fish in a given year), a and b are constants. Maximum sustainable yield (MSY), optimum effort and equilibrium yields were then calculated: $MSY = a^2/4b$; optimum effort = $a/2b$; equilibrium yield = $aE - bE^2$.

The theoretical annual yield (Y) of fish was estimated using a range of models suitable for use with non-temperate data. These were based on various measures including lake surface area (A), morphoedaphic index (MEI; the ratio of total dissolved salts (TDS) to mean lake depth (Z)), primary productivity ($P = \text{gO}_2 \text{ m}^{-2} \text{ d}^{-1}$, $X = \text{gC m}^{-2} \text{ yr}^{-1}$), air temperature (T) and effort (E) as listed below by author:

- (a) Crul (1992): $Y = 8.32 A^{0.92}$ ($Y = \text{t yr}^{-1}$);
- (b) Henderson & Welcomme (1974): $Y = 8.7489 \text{ MEI}^{0.3813}$ ($Y = \text{kg ha}^{-1}$);
- (c) Melack (1976): $\log Y = 0.113 P + 0.91$ ($Y = \text{kg ha}^{-1}$);
- (d) Oglesby (1977): $Y = 0.00038 X^{2.21}$ ($Y = \text{mg dry wt m}^{-2} \text{ yr}^{-1}$);
- (e) Schleisinger & Regier (1982):
 - (i) $\log_{10} \text{MSY} = 0.061 T + 0.043$ ($Y = \text{kg ha}^{-1}$);
 - (ii) $\log_{10} \text{MSY} = 0.044 T + 0.482 \log_{10} \text{MEI} + 0.021$ ($Y = \text{kg ha}^{-1}$);
 - (iii) $\log_{10} Y = 0.051 T + 0.358 E_{\text{Low}} + 0.161 \log_{10} \text{MEI} - 0.383$ ($Y = \text{kg ha}^{-1}$);
 - (iv) $\log_{10} Y = 0.050 T + 0.349 E_{\text{High}} + 0.146 \log_{10} \text{MEI} - 0.367$ ($Y = \text{kg ha}^{-1}$).

In the foregoing equations the following values were used: $A = 163 \text{ km}^2$; $Z = 3.35 \text{ m}$; $\text{TDS} = 231$; $\text{MEI} = 68.95$; $P = 8.98 \text{ gO}_2 \text{ m}^{-2} \text{ d}^{-1}$; $X = 99.16 \text{ gC m}^{-2} \text{ yr}^{-1}$; $T = 21 \text{ }^\circ\text{C}$; $E_{\text{Low}} = 1$; $E_{\text{High}} = 2$. Sources of data for mean values of TDS and primary production were Melack (1979), Harper (1987), Harper (1991) and original (this study) and, for temperature, Litterick et al.

(1979) and Muchiri (1990). Area is for 1991, the year in which the average depth was measured (see below).

In order to determine the above figure for average depth the lake was surveyed during August 1991 using a Lowrance X-15 chart recording echo-sounder with a 20° transducer beam. The soundings were carried out from a small boat driven at constant speed between two defined positions and, using known travel times and distances, the chart recordings were transcribed into depth profiles. Åse et al. (1986) had used a similar echo-sounder in 1983 but had concentrated on the north of the lake. After confirmation of the 1983 findings, the 1991 survey concentrated on previously unmeasured areas.

From 1987, annual collections of bass and tilapia specimens for gut analysis were obtained by gillnetting (11–50 mm bar mesh) in the major habitat types of open water, submerged macrophyte beds, rocky shoreline and the marginal vegetation fringe. Stomachs were removed from representative samples across the size range captured, for each habitat type from each sampling year, and analysed as described by Muchiri (1990) and Hickley et al. (1994).

Results

Lake depth

A depth contour map based on both actual readings from the 1991 echo-sounder traces and interpolation from the map published by Åse et al. (1986) is given in Figure 1. Note that the lake level was approximately 1.75 m lower in 1991 than in 1983. Two examples of depth profiles, from west to east and south to north, are given in Figure 2.

Commercial fish catches

The annual catches for the gillnet fishery are shown in Figure 3. Over the years there have been great fluctuations in both the amount of fish landed (21–1150 t yr^{-1}) and the number of fishing canoes (6–104). Catch per canoe ranged from 0.73 to 92.8 t yr^{-1} . The average species composition of the catch (1987–1998) was *O. leucostictus* 67.6%, *T. zillii* 10.2% and *M. salmoides* 22.2%. The mean sizes (fork length) of fish landed were *O. leucostictus*, 192 mm (S.D. = 23 mm), *T. zillii*, 168 mm (S.D. = 17 mm), and bass, 262 mm (S.D. = 18 mm).

Three separate phases in the development of the fishery were identified:




Figure 4. Regression plots of computed catch per unit effort (CPUE) on effort (E), together with the resulting equilibrium yield curves, for the whole study period alongside separate plots for the three development phases of the fishery. (a) 1962–1998; $CPUE = 35.29 - 0.485E$ ($r^2 = 0.196$) (b) 1962–1973; $CPUE = 78.48 - 1.897E$ ($r^2 = 0.358$) (c) 1974–1988; $CPUE = 15.06 - 0.148E$ ($r^2 = 0.435$) (d) 1990–1998; $CPUE = 2.59 - 0.237E$ ($r^2 = 0.491$).

recent times, the average CPUE (3.5 t yr^{-1}) has been at an all time low and MSY could not be calculated.

Theoretical annual fish yields are presented in Table 1(b). The average estimate shows the potential fishery yield to be 889 t yr^{-1} , more than previously recorded MSY values and considerably higher than the 384 t yr^{-1} for the relatively stable 1974–1988 period. The lowest estimates of theoretical yield equate to the 1974–1988 MSY value whereas estimates based on primary production are twice the value of the MSY for the development phase of the fishery (1962–1973).

The River Malewa fishery

The exploitation of the *B. amphigramma* spawning migration up the River Malewa by fishermen using dip-nets was short-lived. Fishing commenced in 1983, catches peaked at 62.9 t yr^{-1} in 1986, decreased to 26.1 t yr^{-1} in 1987 and thereafter only a small number of fish have been caught.

Feeding habits of fish

Micropterus salmoides

Hickley et al. (1994) reported that, until fish reach a size of 260 mm fork length, bass in Lake Naivasha were almost totally dependent upon free-living invertebrate food organisms. Thereafter, additions to the diet included crayfish, fish and frogs. The most important invertebrate prey species for the smaller bass were the water boatman, *Micronecta scutellaris* (Stal.) and dipteran pupae. For bass in the larger length category, the crayfish, *Procambarus clarkii* (Girard), was the preferred food. *Micropterus salmoides* in Lake Naivasha can, therefore, be classed as a generalized macro-predator, principally feeding on free living animals of a kind most likely to be found in the littoral zones.

Table 1. Estimates of actual and theoretical fish yields calculated for Lake Naivasha: (a) Summary statistics for catch per unit effort (CPUE) on effort (E) and the resultant maximum sustainable yield (MSY) (b) Theoretical yields based on various predictive models. For formulae and data sources see Methods

| (a) | | | | |
|-----------------------------|--|---------------------------------------|---|---------------------------|
| Year | E No. canoes (range) | Average CPUE t canoe ⁻¹ | Optimum E No. canoes | MSY t yr ⁻¹ |
| 1962–1998 | 30 (6–104) | 20.52 | 36.34 | 641 |
| 1962–1973 | 16 (6–39) | 46.88 | 19.63 | 731 |
| 1974–1988 | 45 (6–104) | 8.94 | 50.96 | 384 |
| 1990–1998 | 26 (17–35) | 3.52 | (5.46) | (–7) |
| (b) | | | | |
| Model | Parameters used | | Theoretical annual yield t yr ⁻¹ | |
| Crul (1992) | Surface area | | 902 | |
| Henderson & Welcomme (1974) | Morpho-edaphic index (MEI) | | 716 | |
| Melack (1976) | Gross primary production (g O ₂ m ⁻² d ⁻¹) | | 1371 | |
| Oglesby (1977) | Gross primary production (g Cm ⁻² yr ⁻¹) | | 1599 | |
| Schlesinger & Regier (1982) | Air temperature | | 344 | |
| Schlesinger & Regier (1982) | Air temperature & MEI (for depth <25 m) | | 1105 | |
| Schlesinger & Regier (1982) | Air temperature, MEI & effort (low) | | 358 | |
| Schlesinger & Regier (1982) | Air temperature, MEI & effort (high) | | 727 | |
| Mean of estimates | | | | 889 |

Oreochromis leucostictus

Detritus was found to be the principal component of *O. leucostictus* diet (Muchiri, 1990; Muchiri et al., 1994). It is the most abundant food material available to fish in Lake Naivasha and its importance has been previously noted by Malvestuto (1974) and Siddiqui (1977). Of the other dietary constituents, various algae, especially planktonic forms, were predominant although some chironomid larvae and oligochaete worms were also taken. In terms of feeding category classification, *O. leucostictus*, whilst in the main a detritus feeder, can also be described as a micro-herbivore.

Tilapia zillii

As for *O. leucostictus*, the diet of *Tilapia zillii* was reported as being principally detritus (Muchiri, 1990; Muchiri et al., 1994) but this species consumes a significant amount of macrophyte (mostly *Najas* and *Potamogeton*), rather than algae, and relatively large quantities of insect material. *T. zillii* is, therefore, a de-

tritrus feeder like *O. leucostictus* but with a secondary classification of omnivorous browser.

Barbus amphigramma

Muchiri (1990) indicated that *B. amphigramma* is primarily an insectivore, although zooplankton was an important dietary component.

Poecilia reticulata

As the guppy was introduced in an attempt to control mosquito larvae (Litterick et al., 1979), and has been used elsewhere in the tropics for this purpose (Costa, 1985), it is likely that the fish will take any similarly sized aquatic invertebrates to be found in the littoral zone.

Invertebrate fauna

The benthic fauna of Lake Naivasha comprises principally oligochaetes and chironomid larvae (Clark et al., 1989; Muchiri, 1990; Muchiri et al., 1994) a

typical biota of many tropical lakes (e.g. Burgis et al., 1973; Green, 1979; Marshall, 1982). The worms most commonly found in samples were the tubificids *Limnodrilus hoffmeisteri* (Clarepede) and *Branchiura sowerbyi* (Beddard). The predominant chironomid was *Chironomus formosipennis* Keiffer. Recent work (Raburu, 2002) estimated biomass and production of these benthic organisms to be $4.07 \text{ g dry wt m}^{-2} \text{ yr}^{-1}$ and $8.08 \text{ g dry wt m}^{-2} \text{ yr}^{-1}$, respectively.

The littoral macro-invertebrate fauna shows greater species richness with approximately 50 species being present at any one time (Clark et al., 1989; Muchiri, 1990). By far, the most abundant organism recorded in all habitats, ranging from flooded farmland to open water, was the hemipteran *Micronecta scutellaris* (Stal.) at approximately 400 m^{-3} Clark (1992).

Food web

From the information that they obtained on fish diets, Muchiri et al. (1994) constructed a basic food web. In order to facilitate recognition of any under-utilised resource, they restricted the pathways to only the most important of the feeding relationships. Possible under-utilisation was identified in relation to zooplankton, phytoplankton, fish and, in particular, the benthos.

Discussion

Since commercial fishing started, fish have been taken with gill nets which, in theory, makes regulating the fishery easier than if gears such as seine nets and trawls are used. As gill nets are selective, desired mesh sizes can be imposed and effort controlled by limiting numbers and lengths of nets in addition to regulating the numbers of fishing vessels. Unfortunately, the task of enforcing such regulations is often the difficult part for the fisheries manager. The most serious problems to be addressed are those of ensuring conformity with the gill net specifications and size limits by licensed operators and of curbing illegal fishing by unlicensed fishermen who have no regard for regulations. The latter issue is, however, as much a social concern as a fisheries one and requires a combined effort by the Fisheries Department, riparian interests and other stakeholders in the Naivasha region in order to educate and rehabilitate those people involved. A code of conduct has been drawn up by the Lake Naivasha Fisheries Department (1998) in an attempt to foster re-

sponsible fishing practices and to promote compliance with the legislation (Kenya Fisheries Act – Cap 378).

Although fish caught by unlicensed fishermen are unrecorded there was no loss of information on catches brought to the official landing station. Therefore, in spite of difficulties in obtaining accurate data, it could be considered that the reported commercial catches provide a good representation of the pattern of exploitation. Unfortunately, however, in recent years it has been observed that there has been an increasing tendency for more fish to be traded other than at the official location. It is possible, therefore, that the presumed poor performance of the fishery in the 1990–98 period could in part be due to under-reporting rather than reduced catches. Nonetheless, the fact that the lake level and catch show a very recent upturn (Fig. 3), and that in Figure 4(d) CPUE increases with effort, could indicate that the fishery is about to enter a recovery phase.

Calculations of MSY, although based on a simple version of the Shaefer model (Ricker, 1975; Pauly, 1983), are likely to be reliable. Muchiri et al. (1995) also worked on the fish catches for the 15 year period 1974–1988 inclusive. Using sophisticated catch and effort data analysis software (MRAG, 1992) to fit the data to various models of population growth, MSY estimates of $365\text{--}446 \text{ t yr}^{-1}$ were obtained, as compared with 384 t yr^{-1} given in Table 1.

The theoretical fish yields given in Table 1 were calculated from limnological and primary production data. Hanson & Legget (1982), using a data set of 26 temperate lakes, suggested that total phosphorus and macrobenthos biomass were superior to MEI as predictors of fish yields but reported that for two tropical lakes, George and Naivasha, predictions from total phosphorus fell below observed values. Reasons suggested for this were that for tropical lakes, as against temperate ones, fish production is higher and most commercially important species are herbivores. The same situation of underestimation appears to be true for the use of macrobenthos biomass because when the current estimate of 4.07 g m^{-2} (Raburu, 2002) is used in the various formulae proposed by Hanson & Legget (1982) fish yield estimates of only $70\text{--}225 \text{ t yr}^{-1}$ are obtained. Recently, Welcomme (1999) considered methods for estimating the potential output from fisheries and concluded that there is no system as yet to predict the asymptotic level at which Y_{max} will occur other than the generalised predictors such as MEI. Accordingly, the theoretical fish yields for Lake Naivasha were calculated in such a way.

The potential fish yields, with a mean of 889 t yr⁻¹, suggest that the overall MSY for the Naivasha fishery could be higher. In particular, the estimates based on primary production models (Melack, 1976; Oglesby, 1977) and on the Schlesinger & Regier (1982) model for shallow lakes produced yield estimates (1105–1599 t yr⁻¹) around double the overall MSY of 641 t yr⁻¹. These results suggest that Lake Naivasha may have a potential for a larger fisheries output than is realised at the present time.

Tropical lakes typically support many species of fish, the number being broadly correlated with basin area (Welcomme, 1999). Albeit considerable variance according to altitude and degree of isolation, from figures derived from Vanden Bosch & Bernacsek (1990) for 160 tropical lakes (Number of species = 5.9 lake area (km²)^{0.2684}), a lake such as Naivasha could be expected to contain in the order of 23 species. Such species would be then be likely to undertake a wide range of feeding strategies (Lowe-McConnell, 1987).

By modifying the classification scheme proposed by Matthes (1964), Hickley & Bailey (1987) identified five major feeding categories for common species of fish found in the large, shallow, papyrus-fringed lagoons which form the perennial waters of the Sudd swamps (R.Nile). Whilst not necessarily fully applicable to a lake like Naivasha, such a classification serves to indicate the types of fish that could be expected in a shallow water tropical system. The categories are mud-feeders, microherbivores, macroherbivores, omnivores and carnivores. Carnivores can be further subdivided into zooplanktivores, bottom-feeders, browsers in vegetation, generalised macropredators and piscivores. In Lake Naivasha the tilapias, *Oreochromis leucostictus* and *Tilapia zillii*, with their large uptake of detritus, represent mud-feeders. *O. leucostictus* also partly fulfils the role of microherbivore by taking green algae and diatoms, although this species alone is unlikely to fully exploit phytoplankton production. Of the larger plant material, some is eaten by *T. zillii* but *Procambarus clarkii*, which forms the basis of a commercial crayfish fishery (Lowery & Mendes, 1977), is capable of cropping the macrophytes to a major extent (Harper et al., 1990). Omnivores are represented by *T. zillii*, *B. amphigramma* and the guppy. Although zooplankton featured in the diet of both *B. amphigramma* and juvenile bass, neither species fulfils the role of zooplanktivore to a degree compatible with reports that offshore secondary production is high (Mavuti & Litterick, 1981; Harper,

1984, 1987). The largemouth bass is a generalized macropredator.

It appears, therefore, that in terms of the above classification the Naivasha fish fauna is not sufficiently species rich. Indeed, it would be difficult for any fish population comprising only five species to fully exploit all potential food resources. This view is supported by Moreau (1997) who, taking account of modeling by Mavuti et al. (1996) using ECOPATH (Christensen & Pauly, 1992), concluded that Lake Naivasha is a good example of trophic under-exploitation. As early as the mid 1970s it was suggested (Siddiqui, 1977) that new species should be introduced to exploit the remaining niches. Such a proposal has now been carried forward into the Lake Naivasha Management Plan (LNRA, 1999) with a view to increasing commercial catches and providing local employment opportunities.

Based on prospective feeding guilds and the actual food web, Muchiri et al. (1994) identified four areas in terms of food and space with respect to the potential for stocking additional species of fish. The most convincing case, and that which would best be addressed in the first instance, is the bottom feeder. Given that it is the benthic oligochaetes and chironomid larvae which are under-utilized, one of the species of *Mormyrus* is likely to be a suitable candidate for introduction. With regard to introducing fish to consume more phytoplankton and zooplankton than is currently taken by the existing species, there is scope but the proposal is less robust. Species such as *Limnothrissa* or *Alestes* could be stocked to occupy the open water and take zooplankton but, as these species would not be susceptible to the existing gill net fishery, new fishing methods would have to be authorised. *Heterotis niloticus* (Cuvier) could be used to target the phytoplankton. *H. niloticus* has been introduced to a number of waterbodies and has established self-reproducing populations without seriously threatening the pre-existing fish fauna (Moreau, 1997) although, in Naivasha, the long term decline in *C. papyrus* (Boar et al., 1999) could impact upon the availability of nest sites. With regard to the introduction of a dedicated piscivore, it might be best to leave all the cropping of fish to be carried out not by a top predator but by the activities of the commercial fishermen.

Clearly, the necessary feasibility appraisals of each potential species must address all aspects of suitability or otherwise. The EIFAC code of practice for species introductions (Turner, 1988) recommends taking into account feeding habits, growth rates, reproductive

strategy, environmental conditions required, competition with resident species and the potential implications of the fishery for the new species. Nonetheless, when assessing the balance of risks associated with further introductions into Lake Naivasha, the overall species composition and present ecological nature of the lake is such that any unforeseen detrimental effects of stocking could be considered to be less serious in the long term than if they occurred elsewhere.

Due to the relatively unstable environmental conditions of Lake Naivasha (Harper et al., 1990), there is need for continuous appraisal of the fishery. Catch records need to be of high quality and so adequate enforcement of the fisheries regulations is essential (Lake Naivasha Fisheries Department, 1998). In particular, the key enforcement areas to be addressed are the non-compliance with the minimum (100 mm) mesh size, the trading of undersized fish, the setting of nets closer than 100 m from the shoreline, and the landing of fish other than at the official landing station. The establishment of more landing and recording stations (LNRA, 1999) would comprehensively improve the robustness of the catch statistics. Also, the number and length of gill nets fished would be a better measure of effort than the number of licensed canoes.

Fishery recommendations

It is concluded that, at the present time, the fish population of Lake Naivasha does not fully exploit the available food resources and the fishery is not as productive as it could be. It is recommended that there be acceptance of the proposal to enhance fish production by the introduction of additional species (LNRA, 1999). Potential species should be selected and full feasibility analysis as to their suitability be carried out as soon as is practicable. In addition, consideration should be given to the possibility of re-establishing the *B. amphigramma* of the 1980s. Building fish passes to facilitate spawning migrations might enable the population to recover sufficiently to be exploited once again.

It could be argued that there should be no introductions to a conserved wetland. Even if it were to be considered desirable, it is totally impractical, however, to attempt to reverse the inheritance of past stocking history. The lake cannot be returned to its former status of having only its single endemic fish species. Therefore, given that under certain circumstances social and economic benefits can be considered to be

as important as conservation, the concept of further introductions becomes acceptable. Effort should be made, however, to draw from an African fish list when considering potential candidate species.

Any introductions must be part of an overall management package which should also include:

- Conservation measures based on sound ecology such as refuge areas, fish passes on the river weirs and close times;
- Appropriate legislation and the enforcement thereof, such as minimum mesh sizes and a ban on trading in undersized fish;
- Education to improve awareness of both user groups and those in authority;
- Addressing of associated social issues such as redeployment of the poachers.

If introductions are to be made, then this should be done sooner rather than later, with a phased approach to the associated measures such that any new restrictions are accompanied by increasing availability of fish.

Acknowledgements

This study is part of a wider research project at Naivasha undertaken by the Universities of Leicester, U.K., and of Nairobi, Kenya which since 1987 has been authorised by the Office of the President, Government of Kenya under research permit to Dr D. M. Harper no. OP 13/001/12C 46. The project was funded by the Earthwatch Institute, Boston USA and Oxford, England. The data collection would not have been possible without the assistance of numerous Earthwatch volunteers. We are indebted to the Fisheries Department in Naivasha for access to catch data. Our sincere thanks go to the numerous colleagues in Kenya in particular the staff at the erstwhile Elsamere Conservation Centre, Jill & Angus Simpson. The views expressed are those of the authors and not necessarily those of their parent organisations.

References

- Åse, L. E., K. Sembo & P. Syren, 1986. Studies of Lake Naivasha, Kenya, and its drainage area. Naturgeografiska Institutionen Stockholms Universitet, Forsningsrapport 63: 1–75.
- Boar, R. R., D. M. Harper & C. S. Adams, 1999. Biomass Allocation in *Cyperus papyrus* in a Tropical Wetland, Lake Naivasha, Kenya. *Biotropica* 31: 411–421.

- Burgis, M. J., J. E. P. C. Darlington, I. G. Dunn, G. G. Ganf, J. J. Gwahaba & L. M. McGowan, 1973. The biomass and distribution of organisms in Lake George, Uganda. *Proc. R. Soc. Lond. B*, 184: 271–298.
- Christensen, V. & D. Pauly, 1992. A Guide to the ECOPATH II Software System. version 2. 1. ICLARM Software, 6.
- Clark, F., 1992. A study of a population of *Micronecta scutellaris*. Hemiptera: Corixidae. in Lake Naivasha, Kenya. *Hydrobiologia* 248: 115–124.
- Clark, F. A. Beeby & P. Kirby, 1989. A study of the macro-invertebrates of Lakes Naivasha, Oloidien and Sonachi, Kenya. *Rev. Hydrobiol. Trop.* 22. 10: 21–23.
- Costa, H. H., 1985. Control of *Culex quinquefasciatus* larvae by the larvivorous fish *Poecilia reticulata* in drains and ditches of Colombo. *J. Natl. Aquat. Resour. Agency Sri Lanka*. 32: 24–33.
- Crul, R. C. M., 1992. Models for Estimating Potential Fish Yields of African Inland Waters. FAO Committee for the Inland Fisheries of Africa. CIFA. Occasional Paper, 16: 22 pp.
- Elder, H. Y., D. T. Garrod & P. J. P. Whitehead, 1971. Natural hybrids of the African cichlid fishes *Tilapia spirulus nigra* and *T. leucosticta*, a case of hybrid introgression. *Biol. J. linn. Soc.* 3: 103–146.
- Green, J., 1979. The fauna of Lake Sonfon, Sierra Leone. *J. Zool. Lond.* 187: 113–133.
- Hanson, J. M. & W. C. Legget, 1982. Empirical prediction of fish biomass and yield. *Can. J. Fish. aquat. Sci.* 39: 257–263.
- Harper, D. M., 1984. Recent changes in the ecology of Lake Naivasha, Kenya. *Verh. int. Ver. fur Theor. Ang. Limnol.* 22: 1192–1197.
- Harper, D. M., 1987. The ecology and distribution of the zooplankton in Lakes Naivasha and Oloidien. In Harper, D. M. (ed.), University of Leicester Studies on the Lake Naivasha ecosystem, 1982–1984; Final Report to the Kenya Government: 101–121.
- Harper, D. M., 1991. Primary production in Lake Naivasha, Kenya. *Verh. int. Ver. Limnol.* 24: 1112–1116.
- Harper, D. M., K. M. Mavuti & S. M. Muchiri, 1990. Ecology and management of Lake Naivasha, Kenya, in relation to climatic change, alien species introductions and agricultural development. *Env. Conserv.* 17: 328–336.
- Henderson, H. F. & R. L. Welcomme, 1974. The Relationship of Yield to Morpho-edaphic Index and Number of Fishermen in African Inland Fisheries. UN/FAO CIFA Occ. Pap. No. 1: 19 pp.
- Hickley, P. & R. G. Bailey, 1987. Food and feeding relationships of fish in the Sudd swamps. River Nile, southern Sudan. *J. Fish Biol.* 30: 147–159.
- Hickley, P., E. North, S. M. Muchiri & D. M. Harper, 1994. The diet of largemouth bass, *Micropterus salmoides*, in Lake Naivasha, Kenya. *J. Fish Biol.* 44: 607–619.
- Lake Naivasha Fisheries Department, 1998. Code of Conduct for the Fisheries Sector of Lake Naivasha. In: LNRA 1999, Lake Naivasha Management Plan: 74–77.
- LNRA, 1999. Lake Naivasha Management Plan. Lake Naivasha Riparian Association, P. O. Box 1011 Naivasha, Kenya: 77 pp.
- Litterick, M. R., J. J. Gaudet, J. Kalf & J. M. Melack, 1979. The limnology of an African lake, Lake Naivasha, Kenya. Manuscript prepared for the SIL-UNEP Workshop on Tropical Limnology, Nairobi: 73 pp. Mimeo.
- Lowe-McConnell, R. H., 1987. *Ecological Studies in Tropical Fish Communities*. Cambridge University Press, Cambridge.
- Lowery, R. S. & A. J. Mendes, 1977. *Procambarus clarkii* in Lake Naivasha, Kenya, and its effects on established and potential fisheries. *Aquaculture* 11: 111–121.
- Malvestuto, S., 1974. The Fishes of Lake Naivasha: Their Biology, Exploitation and Management. Mimeographed report to the Director of Fisheries, Government of Kenya, Nairobi, Kenya: 37 pp, 5 figs.
- Marshall, B. E., 1982. The benthic fauna of Lake McIlwain. In Thornton, J. A. (ed.), Lake McIlwain. *Monographiae Biologica* 49: 144–145.
- Matthes, H., 1964. Les poissons du Lac Tumba et de la région d'Ikela. Etude systématique et écologique. *Ann. Mus. R. Afr. Centr. Ser. 8vo. Ser. Zool.* 126: 1–204.
- Mavuti, K. & M. R. Litterick, 1981. Species composition and distribution of zooplankton in a tropical lake, Lake Naivasha, Kenya. *Arch. Hydrobiol.* 93. 1: 52–58.
- Mavuti, K., J. Moreau, J. Munyandorero & P. D. Plisnier, 1996. Analysis of trophic relationships in two shallow equatorial lakes, Lake Naivasha, Kenya and Lake Ihema, Rwanda. *Hydrobiologia* 321: 89–100.
- Melack, J. M., 1976. Primary production and fish yields in tropical lakes. *Trans. Am. Fish. Soc.* 105: 575–580.
- Melack, J. M., 1979. Photosynthetic rates in four tropical African fresh-waters. *Freshwat. Biol.* 9: 444–571.
- Moreau, J., 1997. Impact of Fish Introductions in African Ecosystems: an Evaluation by Using ECOPATH II. In Remane, K. (ed.), African Inland Fisheries, Aquaculture and the Environment. Oxford: Fishing News Books, Blackwell Science Ltd: 326–350.
- MRAG, 1992. Catch-Effort Data Analysis. Unpublished software manual, MRAG Ltd, London and ODA, London.
- Muchiri, S. M., 1990. The Feeding Ecology of Tilapias and the Fishery of Lake Naivasha, Kenya. PhD Thesis unpublished, University of Leicester, U.K.
- Muchiri, S. M., J. B. Hart & D. M. Harper, 1995. The persistence of two introduced species in Lake Naivasha, Kenya, in the face of environmental variability and fishing pressure. In Pitcher, T. J. & J. B. Hart (eds), *Impact of Species Change in African Lakes*. London: Chapman & Hall: 299–319.
- Muchiri, S. M. & P. Hickley, 1991. The fishery of Lake Naivasha, Kenya. In Cowx, I. G. (ed.), *Catch Effort Sampling Strategies: Their Application in Freshwater Fisheries Management*. Oxford: Fishing News Books, Blackwell Scientific Publications: 382–392.
- Muchiri, S. M., P. Hickley, D. M. Harper & E. North, 1994. The potential for enhancing the fishery of Lake Naivasha, Kenya. In Cowx, I. G. (ed.), *Rehabilitation of Freshwater Fisheries*. Oxford: Fishing News Books, Blackwell Scientific Publications: 348–357.
- Oglesby, R. T., 1977. Fish yield as a monitoring parameter and its prediction for lakes. In Alabaster, J. S. (ed.), *Biological Monitoring of Inland Fisheries*. Barking: Applied Science Publishers.
- Pauly, D., 1983. Some Simple Methods of Assessing Fish Stocks. F. A. O. Rome. *Fish. Tech. Pap. No.* 234: 52 pp.
- Raburu, P., K. M. Mavuti, D. M. Harper & F. L. Clark, 2002. Population structure and secondary productivity of *Limnodrilus hoffmeisteri* (Claparede) and *Branchiura sowerbyi* Beddard in the profundal zone of Lake Naivasha, Kenya. *Hydrobiologia* 488 (Dev. Hydrobiol. 168): 153–161.
- Ricker, W. E., 1975. Computation and interpretation of biological statistics of fish populations. *Bull. Fish. Res. Bd Can.* 191: 382 pp.
- Schlesinger, D. A. & H. A. Regier, 1982. Climatic and morphoedaphic indices of fish yields in natural lakes. *Trans. Am. Fish. Soc.* 111: 141–150.
- Siddiqui, A. Q., 1977. Lake Naivasha, Kenya, East Africa. Fishery and its management together with a note on the food habits of fish. *Biol. Conserv.* 12: 217–218.

- Turner, G. E., 1988. Code of Practice and Manual of Procedures for Consideration of Introductions and Transfers of Marine and Freshwater Organisms. FAO European Inland Fisheries Advisory Commission. EIFAC. Occasional Paper, 23: 44 pp.
- Vanden Bosch, J. P. & G. M. Bernacsek, 1990. Source Book for the Inland Fishery Resources of Africa. FAO Committee for the Inland Fisheries of Africa. CIFA. Technical Paper, 18, Vol. 1: 411 pp., Vol. 2: 240 pp., Vol. 3: 219 pp.
- Welcomme, R. L., 1999. A review of a model for quantitative evaluation of exploitation levels in multi-species fisheries. *Fish. Manage. Ecol.* 6: 1–19.